

MODE OF EMPLACEMENT OF SUEVITE AND IMPACT MELT PARTICLES WITHIN THE CHICXULUB PEAK RING. P. Kaskes^{1,2}, L. Ferrière³, S. Chernonozhkin⁴, S. Goderis¹, S. J. de Graaff^{1,2}, T. Déhais^{1,2}, J.-G. Feignon⁵, C. Koeberl⁵, F. Vanhaecke⁴, and Ph. Claeys¹. ¹Analytical-, Environmental- and Geo-Chemistry, Vrije Universiteit Brussel, Brussels, Belgium (pim.kaskes@vub.be), ²Laboratoire G-Time, Université Libre de Bruxelles, Brussels, Belgium, ³Natural History Museum, Vienna, Austria, ⁴Department of Chemistry, Ghent University, Ghent, Belgium, ⁵Department of Lithospheric Research, University of Vienna, Vienna, Austria,

Résumé: L'analyse géochimique et pétrographique à haute résolution de suéвите et de fragments d'impact melt du forage de IODP-ICDP Expedition 364 a révélé une mise en place complexe liée à l'éjection de matériel fondu, à l'entrée de l'eau de mer, à la résurgence de l'océan et aux mouvements de seiches dans la structure d'impact de Chicxulub (Mexique) durant le premier jour après l'impact.

Introduction: Studying the petrography and geochemistry of impact lithologies from within the ~200-km-wide and ~66 Myr old Chicxulub impact structure (Mexico) is crucial to constrain the timing and magnitude of impact-induced processes at 'ground-zero' and it also aids in understanding the formation of large impact structures in general. The peak ring of the Chicxulub impact structure was drilled during the 2016 IODP-ICDP Expedition 364, revealing new insights in the nature and emplacement of proximal impactites [1,2]. In contrast to previous boreholes in the southern part of the crater (e.g., Yucatán-6 and Yaxcopoil-1), the Expedition 364 M0077A drill core from the northern peak ring region recovered the first continuous impact melt rock and suevite (impact-melt-bearing polymict breccia) sequence on top of crystalline basement [1]. We studied in detail the petrography and geochemistry of the ~100-m-thick suevite succession from this drill core to unravel its emplacement and to reconstruct the crater infill history of the first moments (minutes – hours) after the Chicxulub impact event.

Material & methods: A total of 212 suevite and impact melt rock samples between depths ~616.5 and 732 meters below seafloor (mbsf) were analyzed petrographically and geochemically, by SEM-EDS, μ XRF mapping, and whole-rock ICP-OES, XRF, ICP-MS and INAA [3,4]. Phase-specific analyses on the matrix and impact melt particles were carried out with EMPA and LA-ICP-MS. Digital image analysis on μ XRF maps of thin sections was performed to quantify the degree of sorting and to determine modal abundances and particle parameters of clast types [5].

Results: The M0077A suevite is dominated by vitric melt particles, and, to a lesser degree, by microcrystalline melt particles and clasts from the sedimentary target (i.e., primary and reacted carbonates, no evaporites detected) and crystalline target (shocked granitoids and rare gneiss, amphibolite, and dolerite). The suevite sequence is deposited on top of a brecciated impact melt rock unit (Fig. 1A) and is subdivided into three units that are distinct in their petrography (Fig. 1B-D), geochemistry, and sedimentology, i.e., from base to

top: (1) the ~5.6-m-thick non-graded suevite, (2) the ~89-m-thick graded suevite, and (3) the ~3.5-m-thick bedded suevite units. The graded unit shows an upward-increasing sorting and is clearly depleted in CaO (~20 wt%) and enriched in heavy REEs compared to the other two units (~40 wt% CaO). Within the very well-sorted, bedded suevite unit, a rare 1.2 mm wide, silicate, glassy impact spherule is preserved (Fig. 1D), which is geochemically comparable to altered vitric melt particles. All suevite units have isolated Cretaceous planktic foraminifera preserved within their clastic matrix, suggesting that marine processes were responsible for the deposition of the entire sequence.




Fig. 1. M0077A impactite petrography. (A) Brecciated impact melt rock with plagioclase (pl) and clinopyroxene (cpx) microlites (note a LA-ICP-MS pit). (B) Non-graded suevite unit with partly recrystallized planktic foraminifera (F) and vitric melt clasts (VMC). (C) Graded suevite unit with vitric and microcrystalline (MMC) melt clasts. (D) Bedded suevite unit with a rare impact spherule (sph). [(A) SEM-BSE image and (B-D) PPL microphotographs].

Discussion: We interpret that the returning ocean water that reached the northern peak ring (estimated to be < 30 minutes after impact based on a simplified one-dimensional 'dam-break' model [2]) entered through a N-NE gap in the Chicxulub outer rim [2]. This ocean

resurge was relatively poor in rock debris and caused intense quench fragmentation when it interacted with the underlying hot impact melt rock [6], resulting in the formation of the hyaloclastite-like, non-graded suevite unit (Fig. 2.1). In the following hours, the crater was continually flooded by an ocean resurge rich in rock debris, which extinguished phreatomagmatic processes and deposited the ~89-m-thick graded suevite unit (Fig. 2.2). After the energy of this resurge slowly dissipated, oscillating seiche waves took over the sedimentary regime and formed the bedded suevite unit (Fig. 2.3).

The single impact spherule preserved at the base of this unit represents a melt droplet [7] that was quenched and ejected into the proximal Gulf of Mexico region in the first minutes after impact. It was then brought back in the final stage of the resurge into the crater to become part of the first seiches. The upper 20 cm of the bedded suevite unit is cross-bedded and rich in terrestrial biomarkers and charcoal, most probably transported by a reflected turbid tsunami from the nearest landmass in central Mexico. Based on modelling, these reflected tsunami waves have likely re-entered the impact basin within 24 hours after the impact [2], providing an important upper time constraint for the deposition of the Chicxulub suevite sequence. Finally, the overlying transitional layer is a micritic and pyritic siltstone with only rare melt particles. A green marlstone yielding a positive iridium anomaly (~1 ppb) marks the top part of

this unit, indicating atmospheric fallout of the final ultrafine meteoritic matter [8]. This material is estimated to have been deposited up to 20 years after impact (Fig. 2.4) [8].

Conclusions: The M0077A suevite and impact melt sequence from the Chicxulub peak ring preserved a high-resolution record that provides a unique window for elucidating the dynamics and timing of marine cratering processes in the direct aftermath (< 1 day) of a large impact event. Correlating the M0077A record with other Yucatán drill cores will help to test to what extent the model of seawater ingress, ocean resurge, and seiches (Fig. 2) can be extrapolated across the entire Chicxulub crater region.

References: [1] Morgan J.V. et al. (2016) *Science*, 354, 878-882. [2] Gulick S.P.S. et al. (2019) *PNAS*, 116, 19342-19351. [3] Kaskes P. et al. (2022) *GSA Bulletin*, 134, 895-927. [4] de Graaff S.J. et al. (2022) *GSA Bulletin*, 134, 293-315. [5] Kaskes P. et al. (2021) *GSA SP*, 550, 171-206. [6] Osinski G.R. et al. (2020) *Geology*, 28, 108-112. [7] Smit J. (1999), *Ann. Rev. Earth Pl. Sc.*, 27, 75-113. [8] Goderis S. et al. (2021) *Sci. Adv.*, 7(9):eabe3647.

Acknowledgments: We thank FWO (project 11E6621N, G0A6517N), BELSPO, VUB Strategic Research Program, FRS-FNRS, University of Vienna Doctoral School (IK-1045), and NSF (OCE-1737087) for supporting this research.




Fig. 2. Chicxulub M0077A drill core lithostratigraphy following [1,3], showing key moments in the deposition of the different suevite units, as highlighted with arrows on the core stratigraphy and as snapshot panels (#1-4), modified from [3].