

FRIPON and IMPACT projects to pinpoint interplanetary matter in the centimetre - hundred meter range

F. Colas¹, S. Sylla^{1,8}, D. Baratoux⁷, S. Bouley^{3,1}, B. Zanda^{2,1}, P. Vernazza⁴, L. Jorda⁴, J. Gattacceca⁵, M. Birlan¹, J.L. Rault^{1,6}, L. Maquet¹, J. Vaubaillon¹, M. Delcroix⁹,

¹ IMCCE, Observatoire de Paris, Paris, France, colas@imcce.fr, ² IMPMC, zanda@mnhn.fr, ³ GEOPS, sylvain.bouley@gmail.com, ⁴ LAM, OSU Pytheas, ⁵ CEREGE, Aix en Provence, ⁶ International Meteor Organization, ⁷ Géosciences Environnement Toulouse, IRD, ⁸Laboratoire Atomes Laser, Dakar, ⁹ Société Astronomique de France.

Summary :

Improving our knowledge of the interplanetary matter is important to better understand the evolution of the solar system. However, it is mostly undetectable except when it enters the terrestrial atmosphere or when it is large enough to be detected by optical telescopes [1] (Harris et al, 2015). There is in particular a lack of knowledge for particles between 100 m (exhaustive limit of large telescopic surveys) and a few centimeters (typical maximum size for meteor surveys) (fig 2). Fireball observation networks like FRIPON (Fireball Recovery and Interplanetary Observation Network) [2] (Colas et al, 2021) can fill this gap for objects between a few centimeters and a few meters. For larger objects, fall statistics on Earth are too low to determine a precise impact flow. The idea of the IMPACT project is to detect the similar falls of objects from several meters to a hundred meters on Jupiter and Saturn: such events should be more frequent on these more massive planets than on the Earth [3] (Delcroix et al 2015).

FRIPON project: The aim of the FRIPON project is to better constrain the connections between meteorites and asteroids. It is easy to study a meteorite in the laboratory but we cannot tell where it came from, because its orbit is most of the time unknown. On the other hand, we currently have more than 750,000 asteroid orbits with almost no physical information. However, these parameters are crucial for understanding the origin and evolution of the solar system.

The goal of the FRIPON network is to detect large objects that are possible sources of meteorites. We decided to use fisheye cameras to detect fireballs of negative magnitude, allowing us to detect particles larger than a few millimeters. To reach a sufficient statistic we decided to cover the entire French territory (fig 1). We measure an average of 2000 orbits a year and expect to get one meteorite a year.




Figure 1: State of the FRIPON network (100 cameras installed, light blue) and extensions in Europe, PRISMA (Italy, green), MOROI (Romania, orange).

Red dots are for radio stations using GRAVES radar, black for station in installation, yellow for new European networks (UK, Belgium, Germany).

IMPACT project: The size distribution of dust and asteroids are known with relative accuracy only in the immediate vicinity of the Earth (fig 2). It is essential to model this flow precisely to understand the formation of the Solar System, in particular to measure the age of planetary surfaces determined by crater counting. The IMPACT project aims to monitoring the surfaces of giant planets with the best temporal continuity to obtain a good statistic of these impacts. It proposes to coordinate observation campaigns with the 1 meter telescope of the Pic du Midi Observatory by associating amateur and professional astronomers in order to maximize the temporal coverage of the observations.

The IMPACT program is complementary to the one carried out by the University of Bilbao, in particular with the development of the DeTeCt program [3] (Delcroix et al, 2013), aiming to automatically detect possible impact flashes in the observations of Jupiter routinely performed by amateur astronomers.




Figure 2 : Cumulative distribution of Near Earth Asteroids [1](Harris and D'Abramo, 2015)




Figure 3 : Impacts of SL9 comet on Jupiter. On the right image, made with CH₄ narrow band filter, it is possible to see (top) the flash and traces of older impacts at the same latitude [4] (Colas et al, 1995)

IMPACT will not only detect live flashes, it will be especially focused on the detection of impact traces with lifetimes of a few days to several months, such as the impact of comet SL9 in 1994 (fig 3). Our goal is to observe Jupiter twice a month, and thus exhaustively detect all traces that last a few weeks or more. With the image quality of the 1m Pic du Midi telescope, we can detect objects from ten meters to a hundred meters. Beyond this size, Jupiter events will be too rare to measure an impact flux with a good precision.

3. Conclusion: With the combination of the FRIPON and IMPACT programs, we hope to be able to measure the flow of inter-planetary material at the Earth's orbit as well as for Jupiter and Saturn, and obtain a new global model for the whole Solar System. With the addition of meteor observations and the discovery of near-Earth asteroids, we will thus have a global assessment of the matter falling on Earth from a few 1/100 mm to 1000 km. The knowledge of this flow will also yield better estimates of the age of cratered surfaces outside those of the Moon that have been calibrated with sample returns.

References:

- [1] Harris, A. W. and D'Abramo, G. (2015). The population of near-earth asteroids. *Icarus*, 257, 302 – 3
- [2] Colas, F., Zanda, B., Bouley, and 340 co-authors M.: FRIPON: a worldwide network to track incoming meteoroid, *A&A*, 2021
- [3] Delcroix, M. and Hueso, R. (2013). Jovian impact flashes detection with DeTeCt software project. *European Planetary Science Congress*, 8, EPSC2013–812.
- [4] Colas, F., Tiphène, D., Lecacheux, J., Drossart, P., de Batz, B., Pau, S., Rouan, D., and Sèvre, F. (1995). Near-infrared imaging of SL9 impacts on Jupiter from Pic-du-Midi Observatory. *Geophysics Research Letters*, 22.

Acknowledgements: FRIPON was initiated by funding from ANR (grant N.13-BS05-0009-03), carried by the Paris Observatory, Muséum National d'Histoire Naturelle, Paris-Saclay University and Institut Pythéas (LAM-CEREGE). FRIPON data are hosted and processed at Institut Pythéas SIP (Service Informatique Pythéas), and a mirror is hosted at IMCCE (Institut de Mécanique Céleste et de Calcul des Éphémérides / Paris Observatory)